



Flora of South Australia

5th Edition | Edited by Jürgen Kellermann

HYDATELLACEAE¹

J.G. Conran²

Subaquatic caespitose annual or perennial herbs; rootstock minute, not forming a rhizome; leaves linear, sessile, alternate, sheathing, eligulate. **Reproductive** units capitulum-like, scapose, heterogamous, unisexual or co-sexual with male florets scattered among female florets; involucre bracts 2–21, lanceolate, spreading or erect, forming an apparent whorl below the florets; perianth absent; male flower a single anther; female flower a superior, \pm trigonous, 1-ovulate ovary, styles free. **Fruit** a shortly pedicellate, achene or follicle-like, sometimes dehiscent, often with 3 prominent longitudinal ribs; seed perispermous, starchy, more or less lacking endosperm, testa smooth or sculptured.

A monogeneric family of 12 species worldwide with 10 endemic to Australia, mainly in south-western W.A. and one species each in New Zealand and India. A single species recorded in S.A.

Hydatella Diels and *Trithuria* Hook.f. were traditionally regarded as two closely-related genera (recognised on the basis of plant sexuality and fruit dehiscence) within the family Centrolepidaceae and allied to the grasses and sedges, but recent molecular studies have shown that they represent a single genus and are members of the basal angiosperm family Hydatellaceae in the order Nymphaeales (Rudall *et al.* 2007; Saarela *et al.* 2007; Sokoloff *et al.* 2008). Although the family consists of a single genus (*Trithuria*), the family name Hydatellaceae (based on the now synonymous *Hydatella*) has priority.

1. TRITHURIA Hook.f.

Fl. Tasman. 2: 78 (1858).

(Greek *treis*, three; *thyris*, window; referring to the 3-way dehiscence of the fruit of *T. submersa*.)

Hydatella Diels, *Bot. Jahrb. Syst.* 35: 93 (1904).

Description and distribution statement as for family (monogeneric family).

1. ***Trithuria submersa*** Hook.f., *Fl. Tasman.* 2: 79, t. 138A (1858). — **Illustr.:** Rudall *et al.*, *Amer. J. Bot.* 94: 1073–1092, fig. 1, 4, 7 (2007); Sokoloff *et al.*, *Taxon* 57: 179–200, fig. 2–10, 13 (2008).

Annual herb, green but usually becoming reddish; leaves lax, straight, narrow-linear, acute, 1–4 cm long, to 1 mm wide. **Reproductive** units numerous, scapose or rarely sessile; scapes erect, filiform, unbranched, accrescent, to 5 cm high, leafless, glabrous; involucre spreading; bracts 4–6, acuminate, 1-veined, subequal, 3–4 mm long, glabrous; male florets 2–5; anther oblong, c. 0.6 mm long, purple; filament 1–2 mm long, rigid; female florets 10–25, c. 2 mm long, densely packed in a hemispherical cluster; ovary ovoid to trigonous, style apparently absent, stigmatic hairs 3–6. **Fruit** a capsule, obovoid-trigonous, c. 0.6 × 0.3 mm, splitting along the 3 ribs, with 3 panels attached at the apex and opening outwards from the base after falling from the pedicel; seed ovoid, c. 0.4 ×

¹ This work can be cited as: Conran, J.G. (2011). Hydatellaceae (version 1). In: Kellermann, J. (ed.), *Flora of South Australia* (ed. 5). 3 pp. (State Herbarium of South Australia: Adelaide). www.flora.sa.gov.au/ed5

² Australian Centre for Evolutionary Biology & Biodiversity, Benham Building DX 650 312, School of Earth & Environmental Sciences, The University of Adelaide, SA 5005, Australia.

© Department of Environment and Natural Resources, Government of South Australia. ISBN 978-1-922027-00-9 (PDF). Publication date: 26 Oct. 2011.



Government
of South Australia


Department of Environment
and Natural Resources

0.25 mm, rugose, brown. **Trithuria. Pl. 1.**

S.A.: EP, MU, KI, SE; W.A.; N.S.W.; Vic.; Tas. Winter annual on mud of stream margins, seasonal swamps and pools. Flowers: Sep.–Nov.

References

- Rudall, P.J., Sokoloff, D.D., Remizowa, M.V., Conran, J.G., Davis, J.I., Macfarlane, T.D. & Stevenson, D.W. (2007). Morphology of Hydatellaceae, an anomalous aquatic family recently recognized as an early-divergent angiosperm lineage. *Amer. J. Bot.* 94: 1073–1092
- Saarela, J.M., Rai, H.S., Doyle, J.A., Endress, P.K., Mathews, S., Marchant, A.D., Briggs, B.G. & Graham, S.W. (2007). Hydatellaceae identified as a new branch near the base of the angiosperm phylogenetic tree. *Nature* 446: 312–315
- Sokoloff, D.D., Remizowa, M.V., Macfarlane, T.D. & Rudall, P.J. (2008). Classification of the early-divergent angiosperm family Hydatellaceae: one genus instead of two, four new species and sexual dimorphism in dioecious taxa. *Taxon* 57: 179–200



Pl. 1. *Trithuria submersa* in natural habitat. Photos: J.G. Conran.