# FILAMENTOUS RED ALGAE PART VII: red algal "creepers"

(Tribe: Herposiphonieae of the Family: Rhodomelaceae)

This key

Formal classification of algae relies on investigating microscopic reproductive features in detail. Often a complete set of reproductive stages is unavailable in the specimens to be investigated, making identification very difficult if the technical systematic literature is used. Fortunately, some algae grow in specific places and some have recognizable shapes that allow them to be sorted directly into the level of Family, Tribe or Genus and so shortcut a systematic search through intricate and often unavailable reproductive features. The pictured key below uses this *artificial* way of searching for a name. Then you can proceed to the appropriate fact sheets or further keys to refine your identification.

The key generally starts with the large and obvious and then proceeds to the smaller and obscure species.

Limitations Unfortunately, to use this key, microscopic investigation of specimens is essential.

Images used below

Unless acknowledged otherwise, all images have been made from pressed specimens or the extensive slide collection of the algal unit, State Herbarium of S Australia, collections generated by the late Professor Womersley and his workers over some 60 years. Images with dark backgrounds have been taken using phase contrast or interference microscopy to highlight transparent structures. Other images may be stained dark blue.

**Scale** The coin used as a scale is 24 mm or almost 1" across.

Names Names follow Womersley (1998) as this continues to provide the most available and comprehensive descriptions of species. Name

changes in the Website Algaebase have been checked.

### THE HERPOSIPHONIEAE

This is a group of **seven** genera in southern Australia, some easily mistaken for a variety of filamentous Red algae.

Microscopic investigation is necessary to separate the Herposiphonieae from other Tribes in the Family: Rhodomelaceae such as the Lophothalieae and Polysiphonieae.

Special features of the Herposiphonieae -


- many plants lie flat on other algae (they are prostrate and epiphytic) or have a substantial creeping base, features from which the Tribe gets its name
- ends of branches often curve downwards or backwards (like snakes about to strike)
- most importantly, two types of side branches (laterals) arise along main branches (axes):
  - short, unbranched laterals or laterals with limited growth (called determinate laterals) that occur either on one side or alternate on opposite sides of axes
  - continually growing (indeterminate) laterals that may branch a few times or may develop into additional axes
  - a particular sequence of determinate and indeterminate branches occurs along axes (although this is sometimes difficult to detect in densely branched species)

Algal names are from Womersley (2003). Currently there have been no name changes in the Website *Algaebase*.

## FIRST STEP

Ensure you are truly looking at the Herposiphonieae by checking if the specimen to be identified has one of the required sequences of determinate and indeterminate laterals illustrated by the diagrams on the next page.

# HERPOSIPHONIEAE: DIAGRAMMATIC EXAMPLE



- main branch (axis, ax); the central filament (c fil) has been exposed
- segments (seg) of pericentral cells
- determinate branches (2 sets shown 1-3 and i-iii)
- indeterminate branches (2 shown -i br)

# DIFFICULTIES:


Separating members of the Herposiphonieae that have upright branches from some epiphytic and creeping *Polysiphonia* spp (in the Tribe: Polysiphonieae) can be difficult.

#### Features in common with Polysiphonia:


- a central filament of cells ringed with pericentral cells of equal length forming segments along branches
- extremely fine, branched hairs trichoblasts near ends of branches
- sporangia in a single line along modified side branches

## ${\bf Features\ in\ } {\it contrast\ } {\bf to\ } {\it Polysiphonia:}$

- plants are sometimes pale red or orange. Polysiphonia species are often dark red, almost black
- trichoblasts when fresh are coloured, not colourless as in Polysiphonia
- a *sequence* of determinate and indeterminate laterals occurs in the Herposiphonieae, but *not* in *Polysiphonia* where laterals look similar to each other



### HERPOSIPHONIEAE: THE BRANCHING PATTERNS




IF THE SPECIMEN TO BE IDENTIFIED HAS ONE OF THE BRANCHING PATTERNS SHOWN ABOVE, THEN PROCEED TO THE NEXT PAGE FOR THE KEY TO MEMBER SPECIES OF THE HERPOSIPHONIEAE

### KEY TO THE HERPOSIPHONIEAE

- 3a. edges of axes and bases of laterals coated (corticated) with cells additional to the 6-7 pericentral cells seen in surface view.

  Figs 1-5 ...... Herpopteros fallax
- 3b. axes un-corticated, of 8 pericentral cells per segment, only 2 large ones obvious in surface view. Figs 6-10. (next page)









Fig. 4: Herpopteros fallax, wide-spreading laterals, corticated at the base; axis with multi-celled adhesive pads on rhizoids





Figs 2, 3: (above & below) two magnifications of *Herpopteros* fallax attached to the Red alga, *Protokuetzingia* (host, **ho**)



with basal cortication (arrowed) running into the axis.

Insert: cross section, 6 pericentral cells visible on each side of the axis



Separating *Herpopteros* from other members of the Polysiphonieae such as *Pitophycos* and *Diplocladia* can be difficult.

See also the appendix: "Herposiphonieae look-alikes"



Fig. 6: Gredgaria maugeana on a piece of sponge



Fig. 8: Gredgaria maugeana female plant, bulbous cystocarps




Fig. 9: *Gredgaria maugeana* sporangial plant, sporangia (arrowed) in a single line, but swelling the lateral




Fig. 7: *Gredgaria maugeana* branched laterals alternating along a flattened axis lacking cortication






Fig10: Gredgaria maugeana cross section of flattened axis: central filament (c fil), large (I-IV) and small (1-4) pericentral cells in view

4a. laterals appear *paired* along the edges of axes 5a. lateral pairs *undivided* ...... 6. 5b. one member of the lateral pairs is divided, the other is *undivided*. Figs 11-13. ..... Dipterosiphonia dendritica 6a. laterals occur in pairs on one side and then the opposite side of axes. Plants grow on fanshaped blades of the Brown algae Zonaria and Lobophora. Figs 14-16. ...... Ditria expleta 6b. pairs of laterals occur on opposite sides of axes

Figs 17, 18 (next page).





Dipterosiphonia dendritica Branches lying on the host (ho) over which they were creeping







Figs 14, 15: Ditria expleta Two magnifications of plants on the flat blade of the Brown alga host Lobophora variegata









Figs 16: Ditria expleta: pairs of undivided laterals, on opposite sides of an axis, one with sporangia





Fig. 17:Dipterosiphonia australica lying on a Red alga host (ho)

australica

laterals paired

across the axis



7a. plants creep over seagrass stems, with erect laterals producing a dense clothing 2-6 mm high. Figs 19, 20.

.....Dipterosiphonia prorepens

7b. plants taller, with branched axes forming a tufted plant body ......8.

8a. plants open-branched, laterals opposite unbranched, producing a fish-bone (pinnate) appearance. 5-7 unbranched laterals between stubby laterals of indeterminate growth. Figs 21, 22. ..... Herposiphoniella plurisegmenta

8b. plants branched variously. 3 unbranched laterals between stubby laterals of indeterminate growth




Fig. 19: Dipterosiphonia prorepens densely clothing a stem of Amphibolis



Fig. 21: Herposiphoniella plurisegmenta



branching pattern: pairs of unbranched laterals along axes, curved at tips




Fig. 22: Herposiphoniella plurisegmenta A genus with 5-7 unbranched laterals between stubby indeterminate laterals (one set above, marked 1-6)

9a.	plants	s relatively robust, axes usually $> 200 \mu m$
	in wid	th 10.
9b.	plants	s slender, axes < 200 µm in width
	10a.	plants occur in a variety of red colours
		including orange-brown; branching
		irregular; axes up to 600µm wide. Figs.
		23, 24.
		Herposiphonia versicolor
	4.04	




Fig. 23: Herposiphonia versicolor. Colour variation of pressed plants

Fig. 24: Herposiphonia
versicolor
backlit specimen,
orange-brown in
colour,
illustrating few
unbranched
laterals between
branched laterals





Sturt Bay. S. Australia.

Polysiphonia rostrata. Fig. 25: Herposiphonia rostrata
on a blade of seagrass



Fig. 26: *Herposiphonia rostrata* variation in form





Fig. 27: *Herposiphonia rostrata* one-sided branching of determinate laterals bearing sporangia

11a.	delicate, low creeping plants, erect parts of	nly
	a few mm tall	12
11b.	plants with erect parts > 2mm tall	13

12b. rare species, 55m deep, running loosely across the bryozoan *Calpidium*, trichoblasts *absent*, pericentral cells usually 12 per segment, indeterminate laterals apparently *dormant*, alternate along either side of the axis. Fig. 32.






Fig. 28: *Herposiphonia calothrix*Portion of a plant extracted from its attachment to a *Codium* plant)






Fig. 30: Herposiphonia calothrix

Three undivided determinate laterals between stubby branched indeterminate laterals; numerous fine rhizoids from the base of the prostrate axis that is curved backwards at the apex



Difficulties:
separating creeping members of *Herposiphonia* from *Lophosiphonia prostrata* and *Polysiphonia teges* can be
difficult.
See the appendix: "Herposiphonieae look-alikes"



13a. plants lowly, 10-30 mm high, branches end in dense tufts of trichoblasts. On the creeping axes, a repeated sequence of *one long* lateral, *one short* lateral then *no* lateral occurs. Figs 33-36.

13b. axes with *one* or *a pair* of short (determinate) laterals on opposite sides of the axis ...... 14.

14b. erect branches 30-60 mm tall, branching open comb-like (pectinate) branching pattern of determinate-laterals, their segments L/B = about 1 or 2.5




Fig.34: *Tiparraria aurata* . Ends of axes curved back, tipped with short trichoblasts

Fig.36: *Tiparraria aurata*. Bulbous, stalked female cystocarps near ends of axes tipped with mature, hair-like trichoblasts




Fig. 37 Herposiphonia filipendula on the red alga Laurencia filiformis (host, ho)



Fig. 33: Tiparraria aurata on the blade of a seagrass





Fig. 35: Tiparraria aurata. creeping axis with sequence of 1 tall, 1 short and no lateral (0) along the segments of its length






Fig. 38: *Herposiphonia filipendula*: slender, creeping runner with short, undivided erect laterals




Fig. 39: *Herposiphonia monilifera*laterals bearing sporangia in bead-like rows, a feature from which the species gets its name




Fig. 40: *Herposiphonia monilifera*Male plant with spermatangial heads (arrowed)



Herposiphonia pectinella





Fig. 42: *Herposiphonia pectinella*, curved apices, segments initially stubby, but rapidly develop the proportions of L:B >1






Fig.44: *Herposiphonia pectinella* thin laterals; swollen female cystocarps releasing carposporangia

# **SPECIES ILLUSTRATED**


species		author(s)	page
Dipterosiphonia	australica	Womersley	6
	dendritica	(C. Agardh) F. Schmitz	5
	prorepens	Falkenberg	6
Ditria	expleta	Huisman	5
Gredgaria	maugeana	Womersley	4
Herposiphonia	calothrix	(Harvey) Womersley	8
	elegans	Womersley	8
	filipendula	(Harvey ex J. Agardh) Falkenberg ex De Toni	9
	monilifera	(Hooker & Harvey) Falkenberg	10
	pectinella	(Harvey) Falkenberg	10
	rostrata	(Sonder) Reinbold	7
	versicolor	(Hooker & Harvey) Reinbold	7
Herposiphoniella	plurisegmenta	Womersley	6
Herpopteros	fallax	Falkenberg	3
Tiparraria	aurata	(Harvey) Womersley	9

# **REFERENCES**

Algaebase. Listing the world's algae <a href="https://www.algaebase.org/">https://www.algaebase.org/</a> accessed January 2025

Womersley, H.B.S. (2003) The Marine Benthic Algae of southern Australia Part IIID. ABRS, Canberra and the State Herbarium of South Australia

# HERPOSIPHONIEAE LOOK-ALIKES






Dasyclonium spp

Left: D. incisum attached to Laurencia elata (host, ho): feather-like laterals with smaller, comb-like laterals on their inner side (adaxial)

 ${\bf Right:}\ D\ flaccidum$ 







Right, above: L. prostrata creeping over

the blade of the Brown

alga *Lobophora* 

Right, below: detail of branching







Baldock, R.N. (2024). Filamentous Red algae Part VII red algal creepers, 13 pp. Algae Revealed